

**159541 to 159581—Continued**

- 159571. Selection D-42-20 from P. I. No. 127460.
- 159572. Selection D-42-21 from P. I. No. 127460.
- 159573. Selection D-37-33 from P. I. No. 127461.
- 159574. Selection D-41-38 from P. I. No. 131353.
- 159575. Selection D-42-26 from P. I. No. 131353.
- 159576. Selection D-42-27 from P. I. No. 131353.
- 159577. Selection D-42-28 from P. I. No. 131353.
- 159578. Selection D-42-33 from P. I. No. 131353.
- 159579. Selection D-42-36 from P. I. No. 131353.
- 159580. Selection D-42-38 from P. I. No. 131353.
- 159581. Selection D-42-42 from P. I. No. 134697.

**159582. PHYSALIS ALKEKENGII L. Solanaceae.**

From Mexico. Seeds presented by the Dirección General de Agricultura, San Jacinto, Distrito Federal. Received Aug. 11, 1947.

**159583. FLEMINGIA sp. Fabaceae.**

From the Belgian Congo. Seeds collected by C. E. Kellogg, Bureau of Plant Industry, Soils, and Agricultural Engineering, Beltsville, Md. Received Aug. 19, 1947.

Introduced as *Flemingia congensis*, an apparently unpublished name. A tropical shrub 4 to 5 feet high with simple leaves and small reddish flowers in few-flowered clusters.

**159584 to 159586.**

From Florida. Plants growing at the United States Plant Introduction Garden, Coconut Grove. Numbered in August 1947.

**159584. CORDIA sp. Boraginaceae.**

Originally received as seeds from Mexico.

**159585. CASUARINA CUNNINGHAMIANA Miquel. Casuarinaceae.**

From Hubert Buckley, St. Petersburg, Fla.

**159586. (Undetermined.)**

Originally collected in Brazil by J. T. Baldwin, Jr.

**159587 and 159588.**

From Brazil. Seeds collected by J. T. Baldwin, Jr., William and Mary College, Williamsburg, Va. Received Dec. 8, 1943.

**159587. (Undetermined.)**

Small tree. Mato Grosso.

**159588. RANDIA sp. Rubiaceae.****159589 to 159592.**

From Florida. Plants growing at the United States Plant Introduction Garden, Coconut Grove. Numbered in August 1947.

**159589. LONCHOCARPUS LUCIDUS Pittier. Fabaceae.**

Originally received from L. W. Smith, Sarasota, Fla.

**159590. VINCETOXICUM PETIOLARE (Gray) Standl. Asclepiadaceae.**

Originally collected by L. N. Goodding, 140 miles south of Nogales, Sonora, Mexico.

**159591. ASCLEPIAS SUBULATA Decaisne. Asclepiadaceae.**

Originally collected as seeds in Sentinel, Ariz. by L. N. Goodding.